

Testing CP and Time Reversal Symmetries with $\Lambda_b \rightarrow \Lambda V(1-)$ Decays

O. Leitner^a and Z. J. Ajaltouni^b

^aLPNHE, Groupe Théorie, Université P. & M. Curie, 4, Place Jussieu, F-75252, PARIS.

^bLPC/IN2P3-CNRS, Université Blaise Pascal, F-63177, AUBIERE.

In this letter, an overview is given for interesting tests of both CP and Time Reversal symmetries with the beauty baryon Λ_b . Extensive use of the helicity formalism and HQET is done for all calculations. Then, emphasis is put on sophisticated methods like analysis of resonance polarizations and particular angle distributions which can exhibit a clear signal of TR violation.

1. PHYSICAL MOTIVATIONS

Modern Particle Physics is well described by gauge field theories which are built on fundamental principles, the most important ones being Lorentz Invariance, Unitarity and Hermiticity. An immediate consequence of these principles is the famous CPT theorem which stipulates that any physical system and its CPT-conjugate one have identical physical laws. On another side, CPT theorem supposes that, if CP symmetry is violated, Time Reversal (TR) is no longer a good symmetry and this mathematical feature represents only an *indirect violation* of TR symmetry. However some experiments performed recently at CERN and Fermilab have shown a clear signal of *direct* TR violation in the $K^0 \bar{K}^0$ system. Our aim is to demonstrate that search for direct TR violation can be performed at the LHC energies by studying the cascade decays of $\Lambda_b \rightarrow \Lambda V$, $\Lambda \rightarrow p\pi$, $V \rightarrow x^+x^-$, x being a lepton or a pion. Our main tool is constructing *Triple Product Correlations* (TPC) defined by the following relations $\vec{v}_i = \vec{p}_i, \vec{s}_i$, $C_{ijk} = \vec{v}_i \cdot (\vec{v}_j \times \vec{v}_k) \rightarrow TR \rightarrow -C_{ijk}$ which are odd by TR. A good example of TPC is the Transverse Polarization of a spinning resonance coming from Λ_b decays.

2. DYNAMICS OF Λ_b DECAYS

At LHC energies, 10% of the produced $b\bar{b}$ pairs hadronize into beauty baryons $\mathcal{B}_b = \Lambda_b, \Sigma_b, \Xi_b, \dots$ and approximately 90% of $\mathcal{B}_b$ are dominated by Λ_b . Recently many authors have pointed out

the possibility of both testing TR invariance and searching for New Physics in Λ_b three-body decays like $\Lambda_b \rightarrow \text{Baryon } \ell^+ \ell^-$, $\text{Baryon } h^+ h^-$, where $\ell^\pm$ and $h^\pm$ could originate from continuum and/or resonances like J/ψ . The model developed in references [1,2] is based on the same final states but, emphasis is put on measuring *physical observables* built from the two intermediate resonances, Λ and J/ψ or $\rho^0 - \omega$ which come from Λ_b decays. Our method offers some advantages: (i) because of Λ_b weak decays, the intermediate resonances are polarized and some components of their vector-polarization $\vec{P}$ are not invariant by TR; (ii) it is expected that Λ_b produced in p-p collisions is *transversally polarized* like ordinary hyperons in hadron collisions and, this physical property will be exploited in order to find many interesting TPC parameters which are T-odd, see Ref. [2].

2.1. Cascade decays

The initial laboratory frame to which Λ_b polarization is referred is defined like: $\vec{e}_1 = \vec{p}_1/p_1$, $\vec{e}_3 = \frac{\vec{p}_1 \times \vec{p}_b}{|\vec{p}_1 \times \vec{p}_b|}$, $\vec{e}_2 = \vec{e}_3 \times \vec{e}_1$ where $\vec{p}_1$ and $\vec{p}_b$ are respectively the incident proton momentum and the produced Λ_b one.

Λ_b being transversally polarized, its polarization value is given by $\mathcal{P}^{\Lambda_b} = \langle \vec{S}_{\Lambda_b} \cdot \vec{e}_3 \rangle$. Let M_i be the Λ_b spin projection along $\vec{e}_3$ axis and, λ_1 and λ_2 are respectively the helicity values of Λ and V . Conservation of total angular momentum leads to *four* possible values for the pair $(\lambda_1, \lambda_2) = (1/2, 0), (1/2, 1), (-1/2, -1), (-1/2, 0)$

Table 1

Branching ratio, $\mathcal{BR}$, for $\Lambda_b \rightarrow \Lambda J/\Psi$, $\Lambda_b \rightarrow \Lambda \rho^0$ and $\Lambda_b \rightarrow \Lambda \omega$.

N_c^{eff}	2	2.5	3	3.5
$\Lambda J/\psi$	8.95×10^{-4}	2.79×10^{-4}	0.62×10^{-4}	0.03×10^{-4}
$\Lambda \rho^0$	1.62×10^{-7}	1.89×10^{-7}	2.2×10^{-7}	2.4×10^{-7}
$\Lambda \omega$	22.3×10^{-7}	4.75×10^{-7}	0.2×10^{-7}	0.64×10^{-7}

with $M_i = \pm 1/2$, $M_f = \lambda_1 - \lambda_2 = \pm 1/2$. In the following, the Jacob-Wick-Jackson helicity formalism will be associated with the method of successive or cascade decays. We define (i) ρ^{Λ_b} the polarization density matrix (PDM) of Λ_b , (ii) $A_0(M_i) = \mathcal{M}_{\Lambda_b}(\lambda_1, \lambda_2) D_{M_i M_f}^{1/2*}(\phi, \theta, 0)$ the Λ_b decay amplitude, (iii) $A_1(\lambda_1)$ and $A_2(\lambda_2)$ the resonance decay amplitudes. So, the total decay amplitude is expressed by:

$$\mathcal{A}_I = \sum_{\lambda_1, \lambda_2} A_0(M_i) A_1(\lambda_1) A_2(\lambda_2),$$

from which the decay probability can be deduced,

$$d\sigma \propto \sum_{M_i, M_i'} \rho_{M_i M_i'}^{\Lambda_b} \mathcal{A}_I \mathcal{A}_I^*.$$

2.2. Hadronic matrix element (HME)

Full computation of the HME has been performed by using the Operator Product Expansion (OPE) formalism supplemented by hypothesis derived from the Heavy Quark Effective Theory (HQET). In the framework of the factorization hypothesis, the HME can be written as: HME = current products $\otimes$ form factors, where (i) form factors are computed according to an expansion with $\mathcal{O}(1/m_b)$ and, (ii) the currents are evaluated from the effective hamiltonian, $\mathcal{H}^{eff} = \frac{G_F}{\sqrt{2}} V_{qb} V_{qs}^* \sum_{i=1}^2 c_i(m_b) O_i(m_b)$, which includes both the *soft* (non-perturbative) contribution expressed by the operators $O_i(m_b)$ and the *hard* (perturbative) one represented by the Wilson coefficients $c_i(m_b)$. Both tree and penguin diagrams are computed and the final hadronic matrix element is given by:

$$\mathcal{A}(\Lambda_b \rightarrow \Lambda V) = \frac{G_F}{\sqrt{2}} f_V E_V \langle \Lambda | \bar{s} \Gamma_\mu b | \Lambda_b \rangle \left\{ \mathcal{M}_{\Lambda_b}^T(\Lambda_b \rightarrow \Lambda V) - \mathcal{M}_{\Lambda_b}^P(\Lambda_b \rightarrow \Lambda V) \right\},$$

with $\Gamma_\mu = \gamma_\mu(1 - \gamma_5)$, $\mathcal{M}_{\Lambda_b}^{T,P}(\Lambda_b \rightarrow \Lambda V) = V_{ckm}^{T,P} A_V^{T,P}(c_i)$, where f_V is the vector-meson decay constant, $A_V^{T,P}(c_i)$ are tree (T) and penguin (P) amplitudes made of combinations of Wilson coefficients according to the nature of $V(1^-)$, V_{ckm} are the Cabbibo-Kobayashi-Maskawa matrix elements and, finally the $\rho^0 - \omega$ mixing is taken into account for the case $V \rightarrow \pi^+ \pi^-$.

3. MAIN PHYSICAL RESULTS

Several results which can be tested experimentally have been obtained. The most important ones are:

3.1. Branching ratios

In the framework of the factorization hypothesis, the color number is an effective parameter which is let *free*. The different branching ratios are proportionnal to the following width given by:

$$\Gamma(\Lambda_b \rightarrow \Lambda V) = \frac{E_\Lambda + M_\Lambda}{M_{\Lambda_b}} \frac{P_V}{16\pi^2} \int_\Omega |A_0(M_i)|^2 d\Omega,$$

while the only experimental branching ratios (Ref. [3]) is $\mathcal{BR}^{exp}(\Lambda_b \rightarrow \Lambda J/\psi) = (4.7 \pm 2.1 \pm 1.9) \times 10^{-4}$ which permits to state that $2.0 \leq N_c^{eff} \leq 3.0$. Numerical results are shown in Table 1.

3.2. Polarizations and asymmetries

Values of the resonance polarizations as well as their density matrix are essential ingredients for Monte-Carlo simulations, particularly for angular distributions. They can be computed from our model as well as other parameters like Λ helicity asymmetries. In Table 2 are listed the numerical results: It is worth noticing that (i) all these "geometrical" parameters do not depend on N_c^{eff}

Table 2

Polarizations and asymmetries in case of $\Lambda_b \rightarrow \Lambda V$ with V is $\rho^0(\omega)$ or J/Ψ .

Parameter	$\Lambda\rho^0 - \omega$	$\Lambda J/\psi$
$\alpha_{AS}^{\Lambda_b}$	0.194	0.490
$\mathcal{P}^\Lambda$	-0.21	-0.17
ρ_{+-}^Λ	0.31	0.25
ρ_{00}^V	0.79	0.66

but they are related directly to the weak decay process and, (ii) longitudinal polarizations of the vector mesons are *dominant*.

3.3. Effects of $\rho^0 - \omega$ Mixing




Figure 1. Branching ratio asymmetry, $a_{CP}(\omega)$, as a function of the $\pi^+\pi^-$ invariant mass in case of $\Lambda_b \rightarrow \Lambda\rho^0(\omega) \rightarrow \Lambda\pi^+\pi^-$ and for $N_c^{eff} = 3$.

The asymmetry parameter between two conjugate channels defined by:

$$a_{CP}(s_\rho) = \frac{\mathcal{BR}(\Lambda_b) - \mathcal{BR}(\bar{\Lambda}_b)}{\mathcal{BR}(\Lambda_b) + \mathcal{BR}(\bar{\Lambda}_b)},$$

where s_ρ is $\pi^+\pi^-$ invariant mass, varies with N_c^{eff} and it is usually too small, $a_{CP} \leq 10^{-3}$. However in the case of $\rho^0 - \omega \rightarrow \pi^+\pi^-$, this asymmetry is *amplified* in the vicinity of the ω mass [4], as it is shown on Fig. 1, and it reaches 7.5% at

the ω pole for $N_c^{eff} = 3.0$. This process is a new way to detect Direct CP Violation, as it has been already shown in beauty meson B decays.

4. TIME ODD OBSERVABLES

The initial laboratory frame is transposed to the Λ_b rest-frame as $(\vec{e}_X, \vec{e}_Y, \vec{e}_Z)$ with $\vec{e}_Z$ parallel to $\vec{e}_3$. For each resonance with momentum $\vec{p}$, a new frame is defined (Jackson [5]) as follows:

$$\vec{e}_L = \frac{\vec{p}}{p}, \quad \vec{e}_T = \frac{\vec{e}_Z \times \vec{e}_L}{|\vec{e}_Z \times \vec{e}_L|}, \quad \vec{e}_N = \vec{e}_T \times \vec{e}_L.$$

4.1. Vector-polarization

Each vector-polarization $\vec{\mathcal{P}}^{(i)}$ can be expanded on the new basis and its components are studied under Parity and Time-Reversal operations.

$$\vec{\mathcal{P}}^{(i)} = P_L^{(i)} \vec{e}_L + P_N^{(i)} \vec{e}_N + P_T^{(i)} \vec{e}_T,$$

with $P_j^{(i)} = \vec{\mathcal{P}}^{(i)} \cdot \vec{e}_j$ and $j = L, N, T$. It could

Table 3

Vector-polarization under Parity and TR operations.

Observable	Parity	TR
$\vec{s}$	Even	Odd
$\vec{\mathcal{P}}$	Even	Odd
$\vec{e}_Z$	Even	Even
$\vec{e}_L$	Odd	Odd
$\vec{e}_T$	Odd	Odd
$\vec{e}_N$	Even	Even
P_L	Odd	Even
P_T	Odd	Even
P_N	Even	ODD

be noticed that if the normal component P_N is non equal to zero, it would be a clear signal of TR Violation. See Table 3 for TR and parity operations on vector polarization.

4.2. Special angles

$\vec{n}_\Lambda$ and $\vec{n}_V$ are defined respectively as the unit normal vectors to Λ and V decay planes. $\vec{n}_\Lambda = \frac{\vec{p}_p \times \vec{p}_\pi}{|\vec{p}_p \times \vec{p}_\pi|}$, $\vec{n}_V = \frac{\vec{p}_{l+} \times \vec{p}_{l-}}{|\vec{p}_{l+} \times \vec{p}_{l-}|}$, or $\vec{n}_V = \frac{\vec{p}_{h+} \times \vec{p}_{h-}}{|\vec{p}_{h+} \times \vec{p}_{h-}|}$.

Those vectors are even under TR; but, if we compute the cosine and the sine of their azimuthal angles, $\phi_{(n_i)} = \phi_{\vec{n}_\Lambda}$, $\phi_{\vec{n}_V}$, as it was suggested by Seghal [6] and Wolfenstein [7] for the decay $K_L^0 \rightarrow \pi^+\pi^-e^+e^-$, $\vec{u}_i = \frac{\vec{e}_Z \times \vec{n}_i}{|\vec{e}_Z \times \vec{n}_i|}$, $\cos \phi_{(n_i)} = \vec{e}_Y \cdot \vec{u}_i$, $\sin \phi_{(n_i)} = \vec{e}_Z \cdot (\vec{e}_Y \times \vec{u}_i)$, we notice that $\cos \phi_{(n_i)}$ and $\sin \phi_{(n_i)}$ are odd under TR; and these asymmetries depend essentially on the azimuthal angle distributions of the Λ resonance in the Λ_b rest-frame, which analytical expression is given by:

$$d\sigma/d\phi \propto 1 + \frac{\pi}{2} \alpha_{AS} \left(\Re e(\rho_{+-}^{\Lambda_b}) \cos \phi - \Im m(\rho_{+-}^{\Lambda_b}) \sin \phi \right).$$

By choosing conservative values for the non-diagonal elements of the Λ_b polarization density matrix: $\Re e(\rho_{+-}^{\Lambda_b}) = -\Im m(\rho_{+-}^{\Lambda_b}) = \sqrt{2}/2$, the following asymmetries are obtained:

- For $\Lambda_b \rightarrow \Lambda J/\psi$, one obtains:
 $AS(\cos \phi_{\vec{n}_\Lambda}) = 4.3\%$,
 $AS(\sin \phi_{\vec{n}_\Lambda}) = -5.5\%$.
- For $\Lambda_b \rightarrow \Lambda \rho^0(\omega)$, one obtains:
 $AS(\cos \phi_{\vec{n}_\Lambda}) = 2.4\%$,
 $AS(\sin \phi_{\vec{n}_\Lambda}) = -2.7\%$.

But, no asymmetries in $\cos \phi_{\vec{n}_V}$ and $\sin \phi_{\vec{n}_V}$ of the resonances $V(1^-)$ are seen. A realistic explanation for these different asymmetries is suggested: (i) T-Odd or TRV effects appear in processes where already Parity is violated like $\Lambda \rightarrow p\pi^-$, (ii) while for processes as $V(1^-) \rightarrow \ell^+\ell^-$, h^+h^- where Parity is conserved, T-Odd effects are absent.

5. CONCLUSION

Complete calculations of the processes $\Lambda_b \rightarrow \Lambda V(1^-)$ have been performed. On the kinematics side, they are based on the helicity formalism and the use of resonance polarization density matrices. On the dynamics side, sophisticated methods using the OPE formalism and HQET have been developed in the framework of the *factorization hypothesis*. Our model is entirely built in the

framework of the Standard Model. No ingredients coming from "Beyond Standard Model" are introduced. The only unknown parameters are the elements of the polarization density matrix of the produced Λ_b in p-p collisions. By adopting conservative values for these matrix elements and an entirely polarized Λ_b , asymmetries related to Time-Reversal Invariance are observed. So, crucial questions may arise: (i) Is there *any dynamics* behind Time-Reversal Violation? (ii) Is this dynamics related to the CKM mechanism? Whatever are the answers, it is plausible to assert that Time-Reversal Violation or T-odd processes are real challenges for the next LHC experiments.

Acknowledgements

One of us (Z.J.A.) is very grateful to Pr Stefan Narison, chairman of the QCD06 International Conference, for his warm hospitality at Montpellier city and for all the illuminating discussions he got with him.

REFERENCES

1. Z. J. Ajaltouni, E. Conte and O. Leitner, Phys. Lett. B614, 165-173 (2005).
2. O. Leitner, Z. J. Ajaltouni and E. Conte, arXiv:hep-ph/0602043.
3. Particle Data Group, S. Eidelman *et al.*, Phys. Lett. B592, 1 (2004).
4. X. H. Guo, O. Leitner and A. W. Thomas, Phys. Rev. D 63, 056012 (2001).
5. J. Jackson, High Energy Physics, Les Houches lectures, edited by C. DeWitt and M. Jacob (Gordon and Breach, New York, 1966 (1965).
6. L. M. Sehgal and J. van Leusen, Phys. Rev. Lett. 83, 4933 (1999).
7. L. Wolfenstein, Phys. Rev. Lett. 83, 911 (1999).